

-
- Margonska, H. B. & D. L. Szlachetko. 2001. *Alatiliparis* (Orchidaceae: Malaxidinae), a new orchid genus with two species from Sumatra. *Ann. Bot. Fenn.* 38: 77–81.
- & ———. 2004. *Disticholiparis*, a new genus of subtribe Malaxidinae. *Orchidee* (Hamburg). 55: 175–179.
- Ormerod, P. 2007. Orchidaceous additions to the flora of China and Vietnam (II). *Taiwania* 52(4): 307–314.
- Pearce, N. R. & P. J. Cribb. 2002. *Liparis* L. C. Rich. Pp. 195–212 in N. R. Pearce & P. J. Cribb (editors), *The Orchids of Bhutan*. Charlesworth Group, Huddersfield, United Kingdom.
- Pridgeon, A. M., P. J. Cribb, M. W. Chase & F. N. Rasmussen. 2005. *Genera Orchidacearum*, Vol. 4: Epidendroideae (Part One). Oxford University Press, New York.
- Ridley, H. N. 1886. A monograph of the genus *Liparis*. *J. Linn. Soc., Bot.* 22: 244–297.
- Seidenfaden, G. 1976. Orchid Genera in Thailand IV. *Liparis* L. C. Rich. *Dansk Bot. Ark.* 31(1): 5–105.
- . 1992. The orchids of Indochina. *Opera Bot.* 114: 128–145.
- Yang, P. H., X. H. Jin & Z. H. Wu. 2009. *Liparis angustioblunga* sp. nov. (Malaxideae: Orchidaceae) from Shannxi, China. *Nordic J. Bot.* 27: 348–350.

Passiflora madidiana, a New Species of Passifloraceae from Northern Bolivia

Peter M. Jørgensen

Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166-0299, U.S.A.
peter.jorgensen@mobot.org

Leslie Cayola

Missouri Botanical Garden and Herbario Nacional de Bolivia, Instituto de Ecología,
Universidad Mayor de San Andrés, Casilla 10077, La Paz, Bolivia. lcayola@yahoo.com

Alejandro Araujo-Murakami

Missouri Botanical Garden and Herbario Nacional de Bolivia, Instituto de Ecología,
Universidad Mayor de San Andrés, Casilla 10077, La Paz, Bolivia; current address:
Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel Rene Moreno,
Casilla 2489, Santa Cruz, Bolivia. araujomurakami@yahoo.com

ABSTRACT. We describe *Passiflora madidiana* P. Jørg., Cayola & Araujo-Murakami (Passifloraceae) as new from Bolivia. It belongs to *Passiflora* L. subg. *Passiflora* and is probably closely related to *P. crassifolia* Killip, *P. mapiriensis* Harms, *P. actinia* Hook., and *P. jilekii* Wawra. The material was collected during a floristic inventory project of the dry forest within the Madidi National Park. We presume the species to be endemic to these highly seasonal and isolated dry forests that surround the upper part of the Río Tuichi within the Madidi National Park.

RESUMEN. Describimos a *Passiflora madidiana* P. Jørg., Cayola & Araujo-Murakami como una especie nueva, que pertenece a *Passiflora* L. subg. *Passiflora* y es probablemente cercana a *P. crassifolia* Killip, *P. mapiriensis* Harms, *P. actinia* Hook. y *P. jilekii* Wawra. El material fue colectado durante el inventario florístico del bosque seco en el Parque Nacional Madidi. Presumimos que esta especie es endémica de estos bosques secos estacionales y aislados que están en los alrededores de la parte alta del río Tuichi dentro el Parque Nacional Madidi.

Key words: Bolivia, dry forests, IUCN Red List, Madidi National Park, *Passiflora*, Passifloraceae.

As a result of making quantitative inventories, new species occasionally emerge. The new species described here was first collected in 2003, but flowering material was not secured until a concerted effort was made to inventory the dry forest in and around the upper part of Río Tuichi in 2004–2005. Both

inventories were part of the Proyecto Madidi, initiated in 2001, with the aim of making a botanical inventory of the Madidi region in northern Bolivia and educating university students in the process. We describe the new species here in order to make the name available for incorporation in the upcoming Bolivian catalog of vascular plants (Jørgensen et al., in prep.).

Passiflora madidiana P. Jørg., Cayola & Araujo-Murakami, sp. nov. TYPE: Bolivia. La Paz: Franz Tamayo, Parque Nac. y Área Natural de Manejo Integrado Madidi, Sect. Pintata próx. al Río Tuichi entre las comunidades de Virgen del Rosario y Suyu Suyu, bosque seco semideciduo en filos y cimas de cerros, 14°26'34"S, 68°34'47"W, 1150 m, 4 Dec. 2005 (fl., fr.), A. Araujo-Murakami, A. Fernández, S. Paredes, E. Cuevas & C. Cuevas 2599 (holotype, LPB; isotypes, BOLV, MA, MO, US, USZ). Figures 1, 2.

Haec species a *Passiflora crassifolia* Killip caulibus vetustis hirsutis, stipulis bis minoribus, bracteis obtusis adaxialiter glabris atque coronae serie externa ter longiore differt.

Herbaceous to woody vine; stems cylindric, striate, hirsute, older stems dark olive green, younger stems light green, puberulent; stipules (2–)2.4–3(–4.1) × 0.7–1.1(–1.6) cm, reniform, extremely asymmetric, sessile, apex abruptly acuminate, mucronate, base rounded, margin glandular-serrate, hirsute. Petiole (1.5–)2.4–3.2(–3.8) cm, hirsute; glands (2 to)4(to 6), sessile, 1 pair 1–1.5 cm from leaf base, opposite-subopposite, the rest irregularly distributed. Leaf



Figure 1. Fertile habit of *Passiflora madidiana* P. Jørg., Cayola & Araujo-Murakami. Illustration by Carmen Ulloa Ulloa from the holotype Araujo-Murakami *et al.* 2599 (LPB).

blade (5–)7.2–10 × (2.7–)3.4–4.5(–4.9) cm, ovate, entire, rarely irregularly bilobed or trilobed, when bilobed or trilobed the angles between midvein and lateral veins 34°–38°; entire leaves (1 to)3(to 5)-nerved, angle between lateral veins and midvein (20°–)30°(–36°) in 3-nerved leaves, angle (45°–)55°(–60°) in 5-nerved leaves; apex obtuse, base truncate-cuneate, margin barely glandular-serrate toward the base, revolute; adaxial surface hirsute, pubescence denser along the principal veins, abaxial surface densely hirsute. Peduncle (2–)2.8–3.5 cm, longer in fruit to 3.3–4 cm, hirsute; bracts 1–1.3 × 0.7–0.9 cm, ovate-elliptic, adaxial surface glabrous, abaxial surface hirsute along principal vein, light green or occasionally light yellow, caducous before the fruit matures, apex obtuse, base cuneate, margin glandular-serrate, hirsute; pedicel (0.5–)0.7–0.9 cm, hirsute; flower solitary; sepals 21–22 × 6.5–7 mm with a pronounced keel, abaxial surface puberulent; awn ca. 1.5 mm; petals white, ca. 21 × 5 mm; corona in 3 filiform series, outer series ca. 19 mm, white with lower 1/5 purple, base occasionally white; second series 2.8–3 mm, white with purple apex; third series 1–1.5 mm, white with purple apex; operculum (1.5–)2–3.5(–4) mm tall, purple, with a filament series attached on the outside; limen ca. 1 mm, placed at 2/3 of the distance between the base of the androgynophore and the operculum; androgynophore ca. 7.5 mm; free part of filaments ca. 5.5 mm; anthers ca. 6.5 × 2.1 mm; ovary ca. 5 × 3 mm, green; style ca. 7 mm, light green, stigma ca. 2 mm, green. Fruit (3–)3.4–3.6 cm, ovoid, immature green, drying dark brown; seeds 4.5 × 3 mm, foveolate with 20 to 22 pits, dark brown.

phore and the operculum; androgynophore ca. 7.5 mm; free part of filaments ca. 5.5 mm; anthers ca. 6.5 × 2.1 mm; ovary ca. 5 × 3 mm, green; style ca. 7 mm, light green, stigma ca. 2 mm, green. Fruit (3–)3.4–3.6 cm, ovoid, immature green, drying dark brown; seeds 4.5 × 3 mm, foveolate with 20 to 22 pits, dark brown.

Distribution, habitat, and IUCN Red List category. *Passiflora madidiana* is found in semideciduous dry forest, along the Río Tuichi in the Madidi National Park at 850–1150 m elevation. We consider the species to be Vulnerable (VU B1,C) following IUCN Red List criteria (IUCN, 2001; IUCN Standards and Petitions Working Group, 2008). The threat to the species is limited if Madidi National Park is respected, but several places within the dry forest area are at risk of fragmentation as a result of increasing cattle and farming activities. So far *P. madidiana* has been collected in three different settings, first on the upper slopes and ridges where the vegetation is somewhat stunted and dominated by *Tabebuia aurea* (Silva Manso) Benth. & Hook. f. ex S. Moore, *Jacaranda mimosifolia* D. Don, *Hymenaea courbaril* L., *Pseudobombax septenatum* (Jacq.) Dugand, *P. longiflorum* (Mart. & Zucc.) A. Robyns, *Aralia soratensis* Marchal, *Anadenanthera colubrina* (Vell.) Brenan, *Trichilia elegans* A. Juss., *Agonandra excelsa*



Figure 2. Flowering branch of *Passiflora madidiana* P. Jørg., Cayola & Araujo-Murakami, from the type Araujo-Murakami et al. 2599. Photograph by A. Araujo-Murakami.

Griseb., *Astronium urundeuva* (Allemão) Engl., and *A. fraxinifolium* Schott ex Spreng. The new species has also been found in taller semideciduous forest with a canopy height of 15–20 m dominated by *Anadenanthera colubrina*, *Tabebuia impetiginosa* (Mart. ex DC.) Standl., *Phyllostylon rhamnoides* (J. Poiss.) Taub., *Astronium urundeuva*, *Machaerium scleroxylon* Tul., *Acacia riparia* Kunth, *Allophylus edulis* (A. St.-Hil., Cambess. & A. Juss.) Radlk., *Ceiba boliviana* Britten & Baker f., *Ximenia americana* L., and *Zanthoxylum fagara* subsp. *lentiscifolium* (Humb. & Bonpl. ex Willd.) Reynel as emergent and canopy-forming species, with an understory composed of *Trichilia catigua* A. Juss., *Amyris* P. Browne, *Ruprechtia apetala* Wedd., *Maytenus cardenasii* Rusby, and *Urera baccifera* (L.) Gaudich. ex Wedd., and in some places dominated by *Chusquea* Kunth and three species of Cactaceae—*Praecereus euchlorus* (F. A. C. Weber) N. P. Taylor, *Cereus tacuaralensis* Cárdenas, and *Opuntia brasiliensis* (Willd.) Haw. The third

location was a semideciduous dry forest on slopes dominated by *Anadenanthera colubrina*, *Oxandra espintana* (Spruce ex Benth.) Baill., *Phyllostylon rhamnoides*, *Machaerium scleroxylon*, and *Capparis polyantha* Triana & Planch.

Phenology. *Passiflora madidiana* has been found with flowers and fruits from late November to early December and in fruit in February. The new species seems to be synchronized with the pronounced changes in temperature and humidity of the rainy season, which starts with torrential rains in November and lasts until March–April.

Etymology and history of discovery. The new species is named after the Madidi National Park, one of the world's most biodiverse protected areas (WCS Bolivia, 2007). *Passiflora madidiana* was first collected in 2003 by Cayola near Sipia in the Madidi National Park; the material consisted only of two